Compared to England:

Better Similar Worse Lower Similar Higher

denotes data refreshed since last quarter

Theme	Aim	Measure Name	Latest Period	Frequency of Reporting	Measure Type	Period 1	Period 2	Period 3* (Latest Period)	Period 1	Period 2	Period 3 (Latest Period)
Overarching To Increase Healthy Life Expectancy		Healthy Life Expectancy at birth - Male (years)	2021-23	Annual	Years	62.7	62.3	61.5	63.3	62.0	63.3
		Healthy Life Expectancy at birth - Female (years)	2021-23	Annual	Years	63.7	62.8	61.9	63.7	62.0	63.0
	To Increase Healthy Life Expectancy	Healthy Life Expectancy at 65 - Male (years)	2021-23	Annual	Years	10.4	10.2	10.1	10.5	9.9	10.5
		Healthy Life Expectancy at 65 - Female (years)	2021-23	Annual	Years	11.7	11.4	11.2	11.3	10.7	11.4

* Explanation of time periods available in the glossary

								England			
Theme	Aim	Measure Name	Latest Period	Frequency of Reporting	Measure Type	Period 1	Period 2	Period 3 (Latest Period)	Period 1	Period 2	Period 3 (Latest Period)
	To Reduce Infant Mortality	Deaths under 1 year per 1,000 live births	2021-23	Annual	Rate per 1,000 live births	3.9	3.9	4.1	5.2	5.1	5.2
	To Reduce illiant Wortanty	Proportion of New Birth Visits completed within 14 days	2023-24	Annual	%	82.7	79.9	83.0	75.6	78.9	84.2
	Stages of development (School readiness / Attainment levels)	% achieving a good level of development (Early Years)	2023-24	Annual	%	65.2%	67.2%	67.7%	67.5%	68.8%	69.9%
Health in Early Life	Stages of development (School readiness / Attainment levels)	% achieving 9-5 pass in English and Maths (Key Stage 4, end of secondary school)	2023-24	Annual	%	50.0%	45.5%	46.2%	45.5%	40.0%	39.7%
	To Reduce Smoking in Pregnancy	Smoking status at time of delivery (% of maternities with known smoking status)	2023-24	Annual	%	9.1%	8.8%	7.4%	10.1%	9.9%	7.7%
	To Increase Breastfeeding	Breastfeeding prevalence at 6-8 weeks	2023-24	Annual	%	49.3%	49.2%	52.7%	36.8%	39.3%	40.7%

^{**} Unavailable due to no national data collection taking place during Covid-19

Compared to England:

Better Similar Worse Lower Similar Higher

denotes data refreshed since last quarter

		England			Staffordshire						
Theme	Aim	Measure Name	Latest Period	Frequency of Reporting	Measure Type	Period 1	Period 2	Period 3 (Latest Period)	Period 1	Period 2	Period 3 (Latest Period)
	To Reduce Hospital Admissions for Self-Harm	Emergency hospital admissions for intentional self-harm ***	2023-24	Annual	DSR per 100,000	163.7	126.2	117.0	169.9	135.2	137.0
		Suicide rate - All persons	2021-23	Annual (3 year pooled)	DSR per 100,000	10.4	10.3	10.7	12.0	11.1	10.5
	To Reduce the Suicide Rate	Suicide rate - Males	2021-23	Annual (3 year pooled)	DSR per 100,000	16.0	15.8	16.4	16.5	15.4	15.6
		Suicide rate - Females	2021-23	Annual (3 year pooled)	DSR per 100,000	5.1	5.2	5.4	7.7	7.0	5.6
Good Mental Health	Improved mental health in young people	New referrals to secondary mental health services (age < 18 years)	2019-20	Annual	DSR per 100,000	4,834	5,994	6,977	7,060	6,421	8,232
	To reduce emergency admissions with a mental health diagnosis in adults	Emergency acute admissions with a mental health diagnosis adults 20+	2023-24	Annual	DSR per 100,000	4,217	3,925	4,202	4,378	4,159	4,819
	To improve the management of depression in Primary Care	The percentage of patients aged 18 or over with a new diagnosis of depression in the preceding 1 April to 31 March who have been reviewed not earlier than 10 days after and not later than 56 days after the date of diagnosis	2023-24	Annual	%	54.9%	65.9%	69.9%	57.7%	69.6%	73.3%
	To improve the management of depression in Primary Care	GP recorded depression (age 18+) %	2022-23	Annual	%	12.3	12.7	13.2	13.3	13.4	14.2

^{***} Self harm hospital admissions have been revised to use re-based ONS population estimates following the 2021 census

Compared to England:	Better	Similar	Worse	Lower	Similar	Higher
	71					

denotes data refreshed since last quarter

			England			Staffordshire					
Theme	Aim	Measure Name	Latest Period	Frequency of Reporting	Measure Type	Period 1	Period 2	Period 3 (Latest Period)	Period 1	Period 2	Period 3 (Latest Period)
	To Reduce Childhood Overweight and Obesity	Reception prevalence of overweight/obesity	2023-24	Annual	%	22.3	21.3	22.1	25.0	22.4	24.4
To Reduce Childhood Ove	To Reduce Childhood Overweight and Obesity	Year 6 prevalence of overweight/obesity	2023-24	Annual	%	37.8	36.6	35.8	37.8	37.6	36.5
Healthy Weight	To reduce adult overweight & obesity	Percentage of adults (aged 18+) classified as overweight or obese	2023-24	Annual	%	63.8	64.0	64.5	68.2	68.5	65.8
	To reduce the prevalence and complications, and improve the management of Type 2 diabetes	Diabetes Prevalence (17+) %	2023-24	Annual	%	7.3	7.5	7.7	7.7	8.0	8.2

^{****} Value not published for data quality reasons

Compared to England:	Better	Similar	Worse	Lower	Similar	Higher				
	denotes data refreshed since last quarter									

							England			Staffordshire		
Theme	Aim	Measure Name	Latest Period	Frequency of Reporting	Measure Type	Period 1	Period 2	Period 3 (Latest Period)	Period 1	Period 2	Period 3 (Latest Period)	
	To reduce emergency hospital admissions in older people	Dementia: Direct standardised rate of emergency admissions (aged 65 years and over)	2019-20	Annual	DSR per 100,000	3,471	3,480	3,517	3,342	3,863	4,146	
	Reduce the number of Older People in Care Homes	SCC brokered Care Home Clients (aged 65 years and over)* Excludes Mental Health & Learning Disability	Jun-25	Annual	Number of Clients	N/A	N/A	N/A	2,332	2,368	2,360	()
		Percentage of deaths that occur in hospital (85+ yrs)	2023	Annual	%	38.8	39.4	38.6	43.1	43.2	41.5	
Healthy Ageing and Frailty	To reduce deaths in Hospital	Percentage of deaths that occur in hospital (75-84 yrs)	2023	Annual	%	47.5	46.9	46.3	49.8	50.7	49.5	
		Percentage of deaths that occur in hospital (65-74 yrs)	2023	Annual	%	48.6	47.3	47.0	50.6	49.7	49.8	
	To improve the management of dementia	Estimated dementia diagnosis rate (age 65 and over)	2025	Annual	%	63.0	64.8	65.6	66.2	68.0	67.9	()
	To reduce falls in older people	Emergency hospital admissions due to falls in people aged 65 and over *****	2023-24	Annual	DSR per 100,000	2,100	1,933	1,984	2,111	2,025	2,299	

***** Falls hospital admissions have been revised to use re-based ONS population estimates following the 2021 census

Glossary:

DSR	II JIPACTIV STANDARDISAD RATA	Allows for differences in the age structure of populations therefore providing fair comparisons to be made between geographic areas, over time and between sexes.
HLE	Healthy Life Eynectancy	A measure of the average number of years a person would expect to live in good health based on contemporary mortality rates and prevalence of self reported good health.
Period 3 (Latest Period)	Period 3 (Latest Period)	Provides data for the most recent period as per the latest period column of the table.
Period 2	IPERIOR /	Provides data for the period prior to Period 3 (latest period). The frequency of reporting column can be used to ascertain how long ago this was e.g. the previous month, the previous year etc.
Period 1	IPERIOR I	Provides data for the period prior to Period 2. The frequency of reporting column can be used to ascertain how long ago this was e.g. the previous month, the previous year etc.

Measure Definitions:

Measure	Description	Numerator	Denominator	Additional Notes
Healthy Life Expectancy at birth		Number of deaths registered and the weighted prevalence of people reporting good or very good health from the Annual Population Survey.	ONS mid-year population estimates, Annual Population Survey sample weighted to local authority population totals.	-
Healthy Life Expectancy at 65	rates and prevalence of self reported good	Number of deaths for ages 65 years and over and the weighted prevalence of people reporting good or very good health from the Annual Population Survey.		-
Deaths under 1 year per 1,000 live births	, - ,	Number of registered infant deaths aged under 1 year.	Number of live births.	-
Proportion of New Birth Visits (NBVs) completed within 14 days	The proportion of infants receiving a new birth visit (NBV) by a Health Visitor within 14 days of birth.		The total number of infants who turned 30 days old in the time period. (The denominator is 30 days and not 14 days because a new birth visit is valid up to 30 days)	-
% achieving a good level of development (Early Years)	Children are defined as having a good level of development at the end of the Early Years Foundation Stage if they are at the expected level for the 12 Early Learning Goals within the 5 areas of learning relating to: communication and language; personal, social and emotional development; physical development; literacy; and mathematics.		Total number of children	Data based on pupil residency not school attended.
% achieving 9-5 pass in English and Maths (Key Stage 4)	The percentage of pupils achieving grade 5 or above in both English and maths GCSEs. To count for this measure a pupil would have to	Number of pupils achieving 9 to 5 passes in both English and maths	Total number of pupils	Data based on pupil residency not school attended.
Smoking status at time of delivery (% of maternities with known smoking status)	, , ,	delivery.	Number of maternities where smoking status is known.	Smoking in pregnancy is associated with various poor health outcomes including the risk of complications in pregnancy and birth, it also increases the risk of low birth weight, premature birth, stillbirth and sudden infant deaths syndrome. (https://www.nhs.uk/pregnancy/keepingwell/stop-smoking/)
Breastfeeding prevalence at 6-8 weeks	Percentage of infants totally or partially breastfed	Number of infants totally or partially breastfed	Infants due a 6 to 8 week review	Important due to breast milk helping to protect from infections, which may result in fewer visits to hospital. It can also reduce the future risk of sudden infant death syndrome, obesity and cardiovascular disease as an adult. (https://www.nhs.uk/conditions/baby/brea stfeeding-and-bottle-feeding/breastfeeding/benefits/)

Measure	Description	Numerator	Denominator	Additional Notes
Emergency hospital admissions for intentional self-harm	lanes Persons	The number of first finished emergency admission episodes in patients, with a recording of self harm in financial year in which episode ended.	Mid-year Population Estimates: Single year of age and sex for local authorities in England and Wales. Source: Office for National Statistics.	High rates may be due to more self-harm in an area or a lower threshold for hospital admission.
Suicide rate	Age-standardised mortality rate from suicide and injury of undetermined intent per 100,000 population	Number of deaths from suicide and injury of undetermined intent.	Population-years (aggregated populations for the three years) for people of ages 10+ only.	-
New referrals to secondary mental health services (age < 18 years)		Number of referrals opening in the financial year to secondary mental health services	Estimates of resident population, Office for National Statistics (ONS)	Data may not represent the activity being delivered, but may be more indicative of the level of recording
Emergency acute admissions with a mental health diagnosis adults 20+		The number of first finished emergency admission episodes in patients with a recording of mental health in financial year in which	Mid-year Population Estimates: Five year of age and sex for local authorities in England and Wales. Source: Office for National Statistics.	Low rates may be due to low rates of diagnosis or good management in the community of those diagnosed.
The percentage of patients aged 18 or over with a new diagnosis of depression in the preceding 1 April to 31 March who have been reviewed not earlier than 10 days after and not later than 56 days after the date of diagnosis	a new diagnosis of depression in the preceding 1 April to 31 March, who have been reviewed not earlier than 10 days after and not later than 56	Patients aged 18 or over with a new diagnosis of depression in the preceding 1 April to 31 March, who have been reviewed not earlier than 10 days after and not later than 56 days after the date of diagnosis.		-
GP recorded depression (age 18+) %		All patients aged 18 or over, diagnosed on or after 1 April 2006, who have an unresolved record of depression in their patient record.	All patients aged 18 or over registered at a GP Practice in England.	High rates may be due to more depression in an area or improved diagnosis locally.
Reception prevalence of overweight/obesity	monitoring purposes children are classified as overweight (including obesity) if their body mass index (BMI) is on or above the 85th centile of the	Number of children in Reception (aged 4-5 years) with a valid height and weight measured by the National Child Measurement Programme with a	Number of children in Reception (aged 4-5 years) with a valid height and weight measured by the National Child Measurement Programme.	-
Year 6 prevalence of overweight/obesity	population monitoring purposes children are classified as overweight (including obesity) if their body mass index (BMI) is on or above the	Number of children in Year 6 (aged 10-11 years) with a valid height and weight measured by the National Child Measurement Programme with a Body Mass Index classified as overweight or living with obesity.	Number of children in Year 6 (aged 10-11 years) with a valid height and weight measured by the National Child Measurement Programme.	-
Percentage of adults (aged 18+) classified as overweight or obese	Percentage of adults aged 18 and over classified as overweight or obese	Number of adults aged 18+ with a Body Mass Index (BMI) classified as overweight (including obese), calculated from the adjusted height and weight variables. Adults are defined as overweight (including obese) if their BMI is greater than or equal to 25kg/m2	Number of adults aged 18+ with valid height and weight recorded.	-
Diabetes Prevalence (17+) %	The percentage of patients aged 17 or over with diabetes mellitus, as recorded on practice disease registers.		Total number of patients aged 17+ yrs registered with the practice.	Percentage may be high due to an area having more diabetics or due to better local identification/diagnosis. Diabetes diagnosis rates in Staffordshire are high compared to the national average, so the high prevalence of diabetes locally is possibly linked to good identification.

Measure	Description	Numerator	Denominator	Additional Notes
Dementia: Direct standardised rate of emergency admissions (aged 65 years and over)	Directly age standardised rate of emergency inpatient hospital admissions for people with a mention of dementia or alzheimer's in any of the diagnosis code positions (aged 65+) per 100,000 population.		Resident population for ages 65 years+.	-
SCC brokered Care Home Clients (aged 65 years and over)* Excludes MH & LD	Number of people aged 65+ in a care home at	Number of clients 65+, excluding learning disability and mental health clients. Staffordshire County Council brokered places only. Clients are Staffordshire resident based.	N/A	-
Percentage of deaths that occur in hospital		Number of registered deaths where the place of death is recorded as hospital.	Total number of registered deaths.	-
Recorded Dementia Diagnoses (age 65 and over)	characteristics of the population and the age and	Services with an unresolved diagnosis of	Applying the reference rates to the registered population yields the number of people aged 65+ one would expect to have dementia within the subject population	-
Emergency hospital admissions due to falls in people aged 65 and over	Emergency hospital admissions for falls injuries in persons aged 65 and over, directly age standardised rate per 100,000.	Emergency admissions for falls injuries. Age at admission 65 and over.	Local Authority estimates of resident population, Office for National Statistics (ONS)	_