

Better Health Evidence Base

Lichfield

Insight Team

March 2023

Background

- Healthy weight identified as a key priority for Staffordshire in the latest Joint Strategic Needs Assessment (JSNA).
- ‘Better Health Staffordshire’ is the branding for a Whole Systems Approach to tackle the causes of excess weight and promote a healthy weight and active lifestyle.
- This shared evidence base sets out Staffordshire’s current position, drivers of excess weight and areas of focus to inform vision work and future planning.
- To be used alongside professional knowledge and other local intelligence.
- Insights will also contribute to a wider evidence base to inform future decision-making on wider determinants that impact on healthy weight and active lifestyles.

Analysis Approach

- Utilised a range of national and local data sources – National Child Measurement Programme (NCMP), Public Health England Profiles, NHS data (NHS digital), Active Lives Survey and more.
- Underpinned by statistical techniques (age standardisation, 95% confidence intervals). If a prevalence is described as higher it will be statistically significantly higher.
- Supported with resident voice intelligence where appropriate.
- Delivered in collaboration with SCC's Public Health and Children and Families teams.
- Data caveats:
 - Some local NCMP data aggregated into 3 year averages due to small numbers. Due to Covid-19, 2020/21 NCMP data has not been released at Local Authority level.
 - Population data uses the BMI classifications for adults and BMI thresholds for children, as recommended by the National Institute for Health and Care Excellence (NICE).
 - BMI classifications should not be used to describe individuals. Positive and sensitive language is encouraged when communicating with individuals and residents.

Key Headlines

- Although overall Lichfield compares favourably with other areas, 1 in 4 Reception children live with excess weight and 1 in 3 Year 6 children live with excess weight, with high levels in parts of the district.
- Levels of Year 6 excess weight remain lower than national levels but could increase over future years as recent Reception age cohorts reach Year 6.
- Healthy weight challenge for children mainly focussed in parts of North Lichfield and Burntwood with high Reception and Year 6 prevalence in Summerfield & All Saints. However, challenges and opportunities exist across the district and shift over time.
- 2 out of every 3 adults live with excess weight in Lichfield, similar to the national average.
- Wider impact on residents health and on the system – Obesity related long term conditions and hospital admissions higher than national.
- Fruit and vegetable consumption and activity levels in Lichfield are similar or better than national but there is a high density of fast food outlets in Town Centres and in areas of higher deprivation.
- COVID-19 likely to have negatively impacted lifestyle behaviours - Staffordshire's residents reported a mixed impact on healthy lifestyles during the first lockdown.


At Reception age ...

In Lichfield...


Levels of excess weight and obesity in Reception both remain similar to national.



1 in 4 live with excess weight
(similar to national)



1 in 10 live with obesity
(similar to national)




Levels of excess weight and obesity have remained stable over the last five years



Excess weight statistically higher than national in Fazeley (30%) and Summerfield & All Saints (30%).
Obesity statistically higher than national in Curborough (16%) and Summerfield & All Saints (15%).

District comparison



- Newcastle is the only district or borough to have a higher than national prevalence of **obesity**.
- Cannock Chase, East Staffordshire Newcastle and Tamworth have a higher than national prevalence of **excess weight**.

By year six...

In Lichfield...

Whilst obesity and excess weight have increased from Reception, levels are similar to or lower than national.



1 in 3 live with excess weight
(lower than national)



1 in 5 live with obesity
(similar to national)




In line with national trends, year 6 excess weight and obesity increased over the last two years



Excess weight statistically higher than national in Summerfield & All Saints (46%)

Obesity statistically higher than national in Summerfield & All Saints (28%)


District comparison



- Across all districts and boroughs, the prevalence of **obesity** is similar to national.
- Across all districts and boroughs, **excess weight** is similar to national apart from Tamworth (higher) and Lichfield (lower).

Areas of focus


Reception excess weight (2018/19, 2019/20 & 2021/22)



Excess weight statistically higher than national in Fazeley (30%) and Summerfield & All Saints (30%).

Obesity statistically higher than national in Curborough (16%) and Summerfield & All Saints (15%).

Year 6 excess weight (2018/19, 2019/20 & 2021/22)




Excess weight statistically higher than national in Summerfield & All Saints (46%).

Obesity statistically higher than national in Summerfield & All Saints (28%).

How does Lichfield compare?


Excess Weight in Reception (2021/22)



- Reception age excess weight is similar to the statistical neighbour average and ranks 5th out of 16 in its statistical neighbour group.
- To be in line with the CIPFA average, there would need to be an additional 10 children of a healthy weight.

Source: Office for Health Improvement and Disparities. Public health profiles.

Excess Weight in Year 6 (2021/22)




- Year 6 excess weight in Lichfield is similar to the statistical neighbour average and ranks 9th out of 16 in its statistical neighbour group.


Source: Office for Health Improvement and Disparities. Public health profiles.

Higher obesity in urban and deprived areas

Excess Weight by Deprivation Decile - Year 6 (2018/19, 19/20 & 21/22)



Excess weight by Urban/Rural classification (2018/19, 19/20 & 21/22)



- For both Reception and Year 6, the prevalence of obesity in Staffordshire's least deprived areas is almost half that of Staffordshire's most deprived areas.
- Excess weight is also more prevalent in our urban areas than in town fringe and more rural locations.
- Consequently, urban and deprived neighbourhoods within the district are more likely to be areas of concern with regard to excess weight and obesity.

Source: NCMP local dataset and PHE fingertips profiles, IMD 2019 and Urban Rural classification. The Indices of Deprivation are a measure of relative deprivation at a local level across England. The IMD combines information from the seven domains to produce an overall relative measure of deprivation. The domains are combined using the following weights: Income Deprivation (22.5%), Employment Deprivation (22.5%), Education, Skills and Training Deprivation (13.5%), Health Deprivation and Disability (13.5%), Crime (9.3%), Barriers to Housing and Services (9.3%), Living Environment Deprivation (9.3%).


Increasing excess weight into adulthood

In Lichfield...



2 in 3 adults live with excess weight, similar to the national average.

Adult Excess Weight Prevalence (Statistical Neighbours 2020/21)




- Lichfield ranks 6th highest of similar local authorities.

Source: Office for Health Improvement and Disparities. Public health profiles.

Districts comparison

Excess weight in adults aged 18 and over, 2020/21




- Excess weight prevalence is higher than national in Cannock Chase, East Staffordshire, Newcastle, South Staffordshire, Staffordshire Moorlands and Tamworth.
- Obesity prevalence is higher than national in Cannock Chase, Newcastle and Tamworth.

Source: Office for Health Improvement and Disparities. Public health profiles.


Healthy eating

- 51% of adults in Lichfield eat five a day, similar to the Staffordshire and national average.
- Across Staffordshire, the density of fast food outlets is similar to national, but generally higher in areas of deprivation and town centres and as a result, sometimes near secondary schools as shown on the map.
- The density of fast food restaurants is lower than national in Lichfield but higher in Stowe ward.

Proportion of adults eating '5-a-day' on a 'usual day' (2019/20)
Public Health England (based on Active Lives, Sport England)



Fast food outlet density (2017)



Physical activity in Lichfield

Regular physical activity is linked to reduced risk of obesity, reduced risk of illness and improved wellbeing.



4 in 10 Lichfield **children** are physically active for one hour a day, similar to Staffordshire and England (2019/20 – latest data for Lichfield).



1 in 3 Lichfield **children** are active for less than 30 minutes a day, similar to Staffordshire and England (2019/20 – latest data for Lichfield).




7 in 10 Lichfield **adults** are active for more than 150 minutes a week, higher than Staffordshire and England (2020/21).



1 in 6 Lichfield **adults** are active for less than 30 minutes a week, lower than Staffordshire and England (2020/21).

High levels of obesity related conditions

Selected Long Term Conditions (QOF registered 2021/22)




In Lichfield, GP registered prevalence of Hypertension, Asthma, Coronary Heart Disease, and Stroke are higher than national.

Note: Prevalences not age standardised – Lichfield has a greater proportion of over 50s than England. The contribution of obesity to each condition varies.

Source [Obesity - NHS \(www.nhs.uk\)](https://www.nhs.uk).

Self reported long term Musculoskeletal problem (GP Survey 2022)
West Midlands Districts and Unitary Authorities



In Lichfield, the self reported prevalence of Musculoskeletal conditions is similar to England.

Source: Office for Health Improvement and Disparities. Public health profiles.

The rate of **obesity related hospital admissions** in South East Staffordshire and Seisdon Peninsula CCG was higher than England (2019/20). Source: NHS Digital (Hospital Episode Statistics)

Supporting Data Matrix

	Cannock Chase	East Staffordshire	Lichfield	Newcastle-under-Lyme	South Staffordshire	Stafford	Staffordshire Moorlands	Tamworth	Staffordshire	England
Reception Excess Weight Prevalence	26%	26%	23%	27%	22%	24%	24%	26%	25%	22%
Statistical Neighbour Rank	4 out of 16	1 out of 16	5 out of 16	1 out of 16	7 out of 16	1 out of 16	2 out of 16	2 out of 16	2 out of 16	
Reception Obesity Prevalence	11%	11%	10%	13%	10%	10%	10%	11%	11%	10%
Statistical Neighbour Rank	4 out of 16	3 out of 16	6 out of 16	1 out of 16	4 out of 16	3 out of 16	1 out of 16	7 out of 16	3 out of 16	
Year 6 Excess Weight Prevalence	39%	40%	34%	37%	38%	36%	39%	41%	38%	38%
Statistical Neighbour Rank	9 out of 16	1 out of 16	9 out of 16	10 out of 16	2 out of 16	2 out of 16	1 out of 16	6 out of 16	2 out of 16	
Year 6 Obesity Prevalence	26%	23%	22%	23%	22%	22%	23%	26%	23%	23%
Statistical Neighbour Rank	5 out of 16	3 out of 16	7 out of 16	7 out of 16	4 out of 16	2 out of 16	2 out of 16	7 out of 16	2 out of 16	
Adult Excess Weight Prevalence	74%	71%	65%	69%	68%	62%	69%	75%	69%	63%
Statistical Neighbour Rank *	2 out of 16	4 out of 16	6 out of 16	5 out of 16	5 out of 16	12 out of 16	1 out of 16	1 out of 16	2 out of 16	
Adult Obesity Prevalence	36%	29%	27%	34%	21%	25%	28%	35%	29%	25%
Statistical Neighbour Rank *	1 out of 16	9 out of 16	7 out of 16	2 out of 16	14 out of 16	8 out of 16	6 out of 16	2 out of 16	2 out of 16	
Physical Activity in Children	**	**	**	43%	42%	47%	42%	**	47%	47%
Physical Activity in Adults	64%	60%	72%	64%	64%	71%	68%	62%	66%	66%
Five a day consumption	45%	55%	51%	54%	52%	56%	58%	53%	53%	55%
Fast Food Outlets (rate per 100,000)	120	119	64	103	51	76	99	86	90	95
Hypertension prevalence	16%	14%	16%	17%	17%	16%	19%	14%	16%	14%
Diabetes prevalence	8%	8%	7%	8%	8%	7%	8%	8%	8%	7%
Coronary Heart Disease prevalence	4%	3%	4%	3%	4%	3%	4%	3%	4%	3%
Asthma prevalence	6%	6%	7%	7%	6%	6%	7%	7%	7%	6%
Stroke prevalence	2%	2%	2%	2%	2%	2%	3%	2%	2%	2%
Musculoskeletal conditions	22%	18%	18%	23%	22%	20%	21%	25%	21%	18%
Obesity related hospital admissions (rate per 100,000 - CCG values) ***	2,708	4,009	2,354	2,311	2,354	2,216	2,311	2,354	2,595	1,615

Worse than England or Statistical Neighbour Group, **Better than England or Statistical Neighbour Group** (difference calculated using 95% statistical significance)

* Difference to Statistical Neighbour Group not calculated ** recent coverage low in the district / borough *** methodology under review. District / Borough figure based on best fit CCG